

UNCLASSIFIED

Defense Technical Information Center
Compilation Part Notice

ADP013714

TITLE: On the q-Bernstein Polynomials

DISTRIBUTION: Approved for public release, distribution unlimited

This paper is part of the following report:

TITLE: Algorithms For Approximation IV. Proceedings of the 2001
International Symposium

To order the complete compilation report, use: ADA412833

The component part is provided here to allow users access to individually authored sections of proceedings, annals, symposia, etc. However, the component should be considered within the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP013708 thru ADP013761

UNCLASSIFIED

On the q -Bernstein polynomials

Halil Oruç and Necibe Tuncer

*Department of Mathematics, Dokuz Eylül University, Tinaztepe Kampüsü
35160 Buca İzmir, Turkey*

halil.oruc@deu.edu.tr, necibe.tuncer@deu.edu.tr

Abstract

We discuss here recent developments on the convergence of the q -Bernstein polynomials $B_n f$ which replaces the classical Bernstein polynomial with a one parameter family of polynomials. In addition, the convergence of iterates and iterated Boolean sum of q -Bernstein polynomial will be considered. Moreover a q -difference operator $\mathcal{D}_q f$ defined by $\mathcal{D}_q f = f[x, qx]$ is applied to q -Bernstein polynomials. This gives us some results which complement those concerning derivatives of Bernstein polynomials. It is shown that, with the parameter $0 < q \leq 1$, if $\Delta^k f_r \geq 0$ then $\mathcal{D}_q^k B_n f \geq 0$. If f is monotonic so is $\mathcal{D}_q B_n f$. If f is convex then $\mathcal{D}_q^2 B_n f \geq 0$.

1 Introduction

First we begin by introducing some notations to be used. For any fixed real number $q > 0$, the q -integer $[k]$ is defined as

$$[k] = \begin{cases} (1 - q^k)/(1 - q), & q \neq 1, \\ k, & q = 1, \end{cases}$$

for all positive integer k . The term Gaussian coefficient is also used, since they were first studied by Gauss (see Andrews [1]).

Let $p(N, M, n)$ denote the number of partitions of a positive integer n into at most M parts, each less than or equal to N . Then the Gaussian polynomial, $G(N, M, n)$, appears as the generating function

$$G(N, M, n) = \begin{bmatrix} N + M \\ M \end{bmatrix} = \sum_{n \geq 0} p(N, M, n) q^n.$$

Note that $\begin{bmatrix} n \\ k \end{bmatrix}$ defined by

$$\begin{bmatrix} n \\ k \end{bmatrix} = \begin{cases} \frac{[n]!}{[r]![n-k]!}, & n \geq k \geq 0, \\ 0, & \text{otherwise,} \end{cases}$$

where $[n]! = [n][n-1] \cdots [1]$ with $[0]! = 1$, is called Gaussian polynomial (or q -binomial coefficient) since it is a polynomial in q with the degree $(n-k)k$. The q -binomial coefficient

cients satisfy the recurrence relations,

$$\begin{bmatrix} n+1 \\ k \end{bmatrix} = q^{n-k+1} \begin{bmatrix} n \\ k-1 \end{bmatrix} + \begin{bmatrix} n \\ k \end{bmatrix} \quad (1.1)$$

and

$$\begin{bmatrix} n+1 \\ k \end{bmatrix} = \begin{bmatrix} n \\ k-1 \end{bmatrix} + q^k \begin{bmatrix} n \\ k \end{bmatrix}. \quad (1.2)$$

The following Euler identity can be verified using the recurrence relation (1.1) by induction that

$$(1+x)(1+qx)\cdots(1+q^{k-1}x) = \sum_{r=0}^k q^{r(r-1)/2} \begin{bmatrix} k \\ r \end{bmatrix} x^r. \quad (1.3)$$

Phillips [8] introduced a generalization of Bernstein polynomials (q -Bernstein polynomials) in terms of q -integers

$$B_n(f; x) = \sum_{r=0}^n f_r \begin{bmatrix} n \\ r \end{bmatrix} x^r \prod_{s=0}^{n-r-1} (1 - q^s x), \quad (1.4)$$

where $f_r = f\left(\frac{[r]}{[n]}\right)$ and an empty product denotes 1. When $q = 1$ the (1.4) reduces the classical Bernstein polynomials. The $B_n(f; x)$ generalizes many properties of classical Bernstein polynomials. Firstly, generalized Bernstein polynomials satisfy the end point interpolation

$$B_n(f; 0) = f(0), \quad B_n(f; 1) = f(1).$$

Phillips [8] also states the generalization of well known forward difference form (see Davis [3]) of the classical Bernstein polynomials by the following theorem.

Theorem 1.1 *The generalized Bernstein polynomial, defined by (1.4), may be expressed in the q -difference form*

$$B_n(f; x) = \sum_{r=0}^n \begin{bmatrix} n \\ r \end{bmatrix} \Delta^r f_0 x^r \quad (1.5)$$

where $\Delta^r f_i = \Delta^{r-1} f_{i+1} - q^{r-1} \Delta^{r-1} f_i$ for $r \geq 1$ and $\Delta^0 f_i = f_i$.

It is easily verified by induction that q -differences satisfy

$$\Delta^r f_i = \sum_{k=0}^r (-1)^k q^{k(k-1)/2} \begin{bmatrix} r \\ k \end{bmatrix} f_{r+i-k}. \quad (1.6)$$

Using the q -difference form of the q -Bernstein polynomials (1.5), one may show that q -Bernstein polynomials reproduce linear functions, since $B_n(1; x) = 1$; $B_n(x; x) = x$.

2 Convergence

In the discussion of the uniform convergence of the q -Bernstein operator, the Bohman-Korovkin Theorem (see Cheney [2]) is used as in the classical case. The Bohman-Korovkin Theorem states that for a linear monotone operator \mathcal{L}_n , the convergence of

$\mathcal{L}_n f \rightarrow f$ for $f(x) = 1, x, x^2$ is sufficient for the sequence of operators \mathcal{L}_n to have the uniform convergence property $\mathcal{L}_n f \rightarrow f, \forall f \in C[0, 1]$. Observe that the q -Bernstein operator is a *monotone linear operator* for $0 < q \leq 1$. For a fixed value of q with $0 < q < 1$

$$[n] \rightarrow \frac{1}{1-q} \quad \text{as } n \rightarrow \infty.$$

Notice that, since $B_n(x^2; x) = x^2 + \frac{x(1-x)}{[n]}$, $B_n(x^2; x)$ does not converge to x^2 . Phillips [8] studies the uniform convergence of q -Bernstein polynomial.

Theorem 2.1 *Let $q = q_n$ satisfy $0 < q_n < 1$ and let $q_n \rightarrow 1$ as $n \rightarrow \infty$. Then,*

$$B_n(f; x) \rightarrow f(x), \quad \forall f(x) \in C[0, 1].$$

The degree of q -Bernstein approximation to a bounded function on $[0, 1]$ may be described in terms of the *modulus of continuity* with the following theorem.

Theorem 2.2 *If f is bounded on $[0, 1]$ and $B_n f$ denotes the generalized Bernstein operator associated with f defined by (1.4), then*

$$\|f - B_n f\|_\infty \leq \frac{3}{2} \omega(1/[n]^{1/2}).$$

An error estimate for the convergence of q -Bernstein polynomials is given in Phillips [8] by the Voronvskaya type theorem.

Theorem 2.3 *Let f be bounded on $[0, 1]$ and let x_0 be a point of $[0, 1]$ at which $f''(x_0)$ exists. Further, let $q = q_n$ satisfy $0 < q_n < 1$ and let $q_n \rightarrow 1$ as $n \rightarrow \infty$. Then the rate of convergence of the sequence of generalized Bernstein polynomials is governed by*

$$\lim_{n \rightarrow \infty} [n](B_n(f; x_0) - f(x_0)) = \frac{1}{2} x_0(1 - x_0) f''(x_0).$$

It is well known that the classical Bernstein polynomials $B_n f$ provide simultaneous approximation of the function and its derivatives. That is if $f \in C^p[0, 1]$, then

$$\lim_{n \rightarrow \infty} B_n^{(p)}(f; x) = f^{(p)}(x)$$

uniformly on $[0, 1]$. It is worthwhile to examine if this property hold for q -Bernstein polynomials. Phillips [7] proved that the p^{th} derivative of q -Bernstein polynomials converges uniformly on $[0, 1]$ to the p^{th} derivative of f under some restrictions of the parameter q . This property results from the generalization of the following theorem.

Theorem 2.4 *Let $f \in C^1[0, 1]$ and let the sequence (q_n) be chosen so that the sequence (ϵ_n) converges to zero from above faster than $(1/3^n)$, where*

$$\epsilon_n = \frac{n}{1 + q_n + q_n^2 + \dots + q_n^{n-1}} - 1.$$

Then the sequence of derivatives of the generalized Bernstein polynomials, $B'_n f$, converges uniformly on $[0, 1]$ to $f'(x)$.

Up to now the convergence of q -Bernstein polynomials is examined by taking a sequence $q = q_n$ such that $q_n \rightarrow 1$ as $n \rightarrow \infty$. In the recent developments, the convergence

of q -Bernstein polynomials is examined for fixed real q , $0 < q < 1$ and for $q \geq 1$. It is proved in Oruç and Tuncer [6] that for a fixed q , $0 < q < 1$, the uniform convergence holds if and only if f is linear on the interval $[0, 1]$. Moreover, if $q \geq 1$, $B_n f \rightarrow f$ as $n \rightarrow \infty$ if f is a polynomial.

Theorem 2.5 *Let $q \geq 1$ be a fixed real number. Then, for any polynomial p ,*

$$\lim_{n \rightarrow \infty} B_n(p; x) = p(x).$$

For any fixed integer i , the q -Bernstein polynomials of monomials (see Goodman *et.al.* [4]) can be written explicitly as

$$B_n(x^i; x) = \sum_{j=0}^i \lambda_j [n]^{j-i} S_q(i, j) x^j, \quad (2.1)$$

where

$$\lambda_j = \prod_{r=0}^{j-1} \left(1 - \frac{[r]}{[n]} \right),$$

an empty product denotes 1, and

$$S_q(i, j) = \frac{1}{[j]! q^{j(j-1)/2}} \sum_{r=0}^j (-1)^r q^{r(r-1)/2} \begin{bmatrix} j \\ r \end{bmatrix} [j-r]^i, \quad 0 \leq i \leq j, \quad (2.2)$$

is the Stirling polynomial of second kind. Thus for any polynomial p of degree m , one may write

$$B_n(p; x) = \mathbf{a}^T \mathbf{A} \mathbf{x}, \quad (2.3)$$

where \mathbf{a} is the vector whose elements are the coefficients of p , \mathbf{A} is an $(m+1) \times (m+1)$ lower triangular matrix with the elements

$$a_{i,j} = \begin{cases} \lambda_j [n]^{j-i} S_q(i, j), & 0 \leq j \leq i, \\ 0, & i < j, \end{cases} \quad (2.4)$$

and \mathbf{x} is the vector whose elements form the standard basis for the space of polynomials P_m of degree m .

Lemma 2.1 *Let $0 < q < 1$ be a fixed real number. Then*

$$\lim_{n \rightarrow \infty} B_n(p; x) = p(x)$$

if and only if $p(x)$ is linear.

This lemma can be generalized for any function $f \in C[0, 1]$.

Theorem 2.6 *Let $0 < q < 1$ be a fixed real number and $f \in C[0, 1]$. Then*

$$\lim_{n \rightarrow \infty} B_n(f; x) = f(x)$$

if and only if $f(x)$ is linear.

3 The iterates

The iterates of classical Bernstein polynomials were first studied by Kelisky and Rivlin [5]. The authors proved that iterates of Bernstein polynomials converge to linear end point interpolants on $[0, 1]$. Several generalization of the result due to Kelisky and Rivlin has been considered by many authors; see Sevy [9] and Wenz [10]. The recent result is the convergence of iterates of generalized Bernstein polynomials. It is proved in Oruç and Tuncer [6] that the q -Bernstein polynomials do preserve the convergence property of iterates of classical Bernstein polynomial. The iterates of generalized Bernstein polynomial are defined by

$$B_n^{M+1}(f; x) = B_n(B_n^M(f; x); x), \quad M = 1, 2, \dots, \quad (3.1)$$

where $B_n^1(f; x) = B_n(f; x)$.

Theorem 3.1 *Let $q \geq 0$ be a fixed real number. Then*

$$\lim_{M \rightarrow \infty} B_n^M(f; x) = f(0) + (f(1) - f(0))x. \quad (3.2)$$

Let A and B be operators then the Boolean sum of A and B is defined to be

$$A \oplus B = A + B - A \circ B.$$

We will be concerned with iterated Boolean sums of the generalized Bernstein polynomials in the form $B_n \oplus B_n \oplus \dots \oplus B_n$ and will denote such an M -fold Boolean sum of the generalized Bernstein operators by $\oplus^M B_n$. Sevy [9] and Wenz [10] proved that the limit of iterated Boolean sums of Bernstein polynomials is the interpolation polynomial with respect to the nodes $(\frac{i}{n}, f(\frac{i}{n}))$ $i = 0, \dots, n$ as $M \rightarrow \infty$. The second theorem of this section will give a result for the convergence of iterates of Boolean sums of generalized Bernstein polynomials. It is proved in Oruç and Tuncer [6] that the iterates of Boolean sums of q -Bernstein polynomials converge to the interpolating polynomial at the nodes $(\frac{[i]}{[n]}, f(\frac{[i]}{[n]}))$.

Theorem 3.2 *The iterated Boolean sum of the q -Bernstein operator $\oplus^M B_n(f; x)$ associated with the function $f(x) \in C[0, 1]$ converges to the interpolating polynomial $L_n f$ of degree n of $f(x)$ at the points $x_i = [i]/[n]$, $i = 0, 1, \dots, n$.*

4 A difference operator \mathcal{D}_q on generalized Bernstein polynomials

Given any function $f(x)$ and $q \in R$ we define the operator \mathcal{D}_q

$$\mathcal{D}_q f(x) = \frac{f(qx) - f(x)}{qx - x}. \quad (4.1)$$

Thus $\mathcal{D}_q f(x)$ is simply a divided difference, $\mathcal{D}_q f(x) = f[x, qx]$. Note that, for a function f and non-negative integer k

$$f[x, qx, \dots, q^k x] = \frac{1}{[k]!} \mathcal{D}_q^k f(x).$$

Theorem 4.1 For any integer $0 \leq k \leq n$,

$$\mathcal{D}_q^k B_n(f; x) = [n] \cdots [n - k + 1] \sum_{r=0}^{n-k} \Delta^k f_r \begin{bmatrix} n-k \\ r \end{bmatrix} x^r \prod_{s=k}^{n-r-1} (1 - q^s x).$$

Proof: Recall the q -difference form of generalized Bernstein polynomials (1.5) and apply the operator \mathcal{D}_q to $B_n(f; x)$ repeatedly k times to get,

$$\mathcal{D}_q^k B_n(f; x) = \sum_{r=0}^{n-k} \frac{[n]!}{[n-k-r]![r]!} \Delta^{k+r} f_0 x^r. \quad (4.2)$$

It will be useful to express Δ^{k+r} in terms of Δ^k . One may prove by induction on m that, for $0 \leq m \leq n - k$ we may write

$$\Delta^{m+k} f_i = \sum_{t=0}^m (-1)^t q^{t(t+2t-1)/2} \begin{bmatrix} m \\ t \end{bmatrix} \Delta^k f_{m+i-t}.$$

Now applying the latter identity to (4.2) gives

$$\mathcal{D}_q^k B_n(f; x) = \sum_{r=0}^{n-k} \sum_{t=0}^r (-1)^t q^{t(t+2k-1)/2} \frac{[n]!}{[n-k-r]![r]!} \begin{bmatrix} r \\ t \end{bmatrix} \Delta^k f_{r-t} x^r. \quad (4.3)$$

Writing $m = r - t$

$$\frac{[n]!}{[n-k-m-t]![m+t]!} \begin{bmatrix} m+t \\ t \end{bmatrix} = \frac{[n]!}{[n-k-m]![m]!} \begin{bmatrix} n-k-m \\ t \end{bmatrix} \quad (4.4)$$

and putting (4.4) in (4.3) we obtain

$$\mathcal{D}_q^k B_n(f; x) = \sum_{m=0}^{n-k} \frac{[n]!}{[n-k-m]![m]!} \Delta^k f_m x^m \sum_{t=0}^{n-k-m} (-1)^t q^{t(t+2k-1)/2} \begin{bmatrix} n-k-m \\ t \end{bmatrix} x^t.$$

Now, it can be easily derived from generalized binomial expansion (1.3), on replacing x by $q^k x$, that

$$\prod_{t=k}^{n-m-1} (1 - q^t x) = \sum_{t=0}^{n-k-m} (-1)^t q^{t(t+2k-1)/2} \begin{bmatrix} n-k-m \\ t \end{bmatrix} x^t.$$

This completes the proof. \square

From Theorem 4.1 we see that, with $0 < q \leq 1$, if $\Delta^k f_r \geq 0$ for $0 \leq r \leq n - k$ then $\mathcal{D}_q^k B_n(f; x) \geq 0$. If f is convex on $0 \leq x \leq 1$ then $\mathcal{D}_q^2 B_n(f; x) \geq 0$ for $0 < q \leq 1$. If f is increasing then $\mathcal{D}_q B_n(f; x) \geq 0$, for $0 < q \leq 1$.

Acknowledgment: The second author is supported from the Institute of Natural and Applied Sciences of D.E.U. and this research is partially supported by the grant AFS 0922.20.01.02.

Bibliography

1. G. E. Andrews, *The Theory of Partitions*, Cambridge University Press, Cambridge, 1998.
2. E. W. Cheney, *Introduction to Approximation Theory*, AMS Chelsea, Providence, 1981.
3. P. J. Davis, *Interpolation and Approximation*, Dover Publications, New York, 1975.
4. T. N. T. Goodman, H. Oruç, and G. M. Phillips, Convexity and generalized Bernstein polynomials, *Proc. Edin. Math. Soc.* **42** (1999) 179–190.
5. R. P. Kelisky and T. J. Rivlin, Iterates of Bernstein polynomials, *Pacific J. Math.* **21** (1967), 511–520.
6. H. Oruç and N. Tuncer, On the convergence and iterates of q -Bernstein polynomials, *J. Approx. Theory*, to appear.
7. G. M. Phillips On generalized Bernstein polynomials, *Numerical Analysis*, D. Griffiths and G. Watson eds. (1996), 263–269.
8. G. M. Phillips, Bernstein polynomials based on the q -integers, The heritage of P. L. Chebyshev: a Festschrift in honor of the 70th birthday of T. J. Rivlin. *Ann. Numer. Math.* **4** (1997), 511–518.
9. J. C. Sevy, Lagrange and least-square polynomials as limits of linear combinations of iterates of Bernstein and Durrmeyer polynomials, *J. Approx. Theory* **80** (1995), 267–271.
10. H. J. Wenz, On the limits of (Linear combinations of) iterates of linear operators, *J. Approx. Theory* **89** (1997), 219–237.